

Lithological control on the spatial evolution of fault slip on the Longitudinal Valley Fault, Taiwan - Supplementary materials

May 23, 2014

S.1 Seismicity

The Longitudinal Valley Fault (LVF) is known to creep near the surface (Angelier et al., 1997; Lee et al., 1998, 2000, 2001, 2005; Chang et al., 2009; Peyret et al., 2011; Champenois et al., 2012) but has also produced large earthquakes, with $M_w > 6.8$ events in 1938, 1951 and 2003 (Fig. S.1). The 2003 Chankung earthquake and the 1951 earthquake sequence are of particular interest since field studies and geodetic analysis have demonstrated that both events have ruptured the aseismic section of the LVF (e.g., Wu et al., 2006; Shyu et al., 2007; Chung et al., 2008; Lee et al., 2008; Mozziconacci et al., 2009; Hsu et al., 2009; Thomas et al., 2014). Those studies have also shown that no significant slip near the surface have been inferred, where the fault is creeping, which supports the hypothesis that the partitioning between aseismic and seismic slip influence the spatial extent of earthquake ruptures (e.g., Kaneko et al., 2010).

Based on the background seismicity, Chung et al. (2008) have suggested that the LVF can be divided into three zones from north to south (Fig. S.1). In the northern section numerous M5 and M6 events have been recorded, with a mix of normal, strike-slip and thrust focal mechanisms, which are likely related to the Ryukyu subduction (Kuoehen et al., 2004). The central zone display a seismic gap with only few earthquakes occurring at depth less than ~ 20 km. The southernmost section on the contrary records more events, predominantly with thrust focal mechanisms.




Figure S1: (a) Regional tectonic setting of the Longitudinal Valley Fault. (b) Seismicity around the LVF from 1991 to 2010. The selected region is based on the assumption that the LVF has a global strike of the fault ($N20^\circ$), is dipping eastward and extending downward to a depth of 30 km. Blue dots represents the $M_w > 3$ seismic events recorded since the Chengkung earthquake (12/10/2003) until December 2010. Black dots represent the $M_w > 3$ seismicity recorded from January 1991 to the day before the Chengkung earthquake (Wu et al., 2008). Yellow, green, red and purple stars indicate the epicenter of the 1908, 1938 and 2003 earthquakes, and the 1951 earthquakes sequence, respectively (Chung et al., 2008). Shyu et al. (2007) estimate of the extent of the November 1951 ruptures is shown in red (the dashed line are inferred).




Figure S2: Geological map of eastern Taiwan (modified from Y. Wang and W.S Chen, 1993) and the locations of samples collected in the field. Circles are samples collected in September 2012, while triangles give the locations for the April 2010 sampling survey. Color attributions for samples and lithological formations are identical.




Figure S3: Representative EDS traces of grains inside the fault gouge, sample lvf4 (Lichi Mélange). For location of the analysis, see Figure 3.3e.

Lichi Mélange - core




Figure S4: Representative EDS traces of grains inside the fault gouge, sample 12w46 (Lichi Mélange). For location of the analysis, see Figure 3.3h




Figure S5: Representative EDS traces of grains inside the Fanshuliao formation, sample T12. For location of the analysis, see Figure 3.2d.

References

- Angelier, J., Chu, H.T., Lee, J.C., 1997. Shear concentration in a collision zone: kinematics of the chihshang fault as revealed by outcrop-scale quantification of active faulting, longitudinal valley, eastern taiwan. *Tectonophysics* 274, 117–143.
- Champenois, J., Fruneau, B., Pathier, E., Deffontaines, B., Lin, K.C., Hu, J.C., 2012. Monitoring of active tectonic deformations in the longitudinal valley (eastern taiwan) using persistent scatterer insar method with alos palsar data. *Earth and Planetary Science Letters* 337, 144–155.
- Chang, S.H., Wang, W.H., Lee, J.C., 2009. Modelling temporal variation of surface creep on the chihshang fault in eastern taiwan with velocity-strengthening friction. *Geophysical Journal International* 176, 601–613.
- Chung, L.H., Chen, Y.G., Wu, Y.M., Shyu, J.B.H., Kuo, Y.T., Lin, Y.N.N., 2008. Seismogenic faults along the major suture of the plate boundary deduced by dislocation modeling of coseismic displacements of the 1951 m7.3 hualien-taitung earthquake sequence in eastern taiwan. *Earth and Planetary Science Letters* 269, 415–425.
- Hsu, Y.J., Avouac, J.P., Yu, S.B., Chang, C.H., Wu, Y.M., Woessner, J., 2009. Spatio-temporal slip, and stress level on the faults within the western foothills of taiwan: Implications for fault frictional properties. *Pure and Applied Geophysics* 166, 1853–1884.
- Kaneko, Y., Avouac, J.P., Lapusta, N., 2010. Towards inferring earthquake patterns from geodetic observations of interseismic coupling. *Nature Geoscience* advance online publication.
- Kuochen, H., Wu, Y.M., Chang, C.H., Hu, J.C., Chen, W.S., 2004. Relocation of eastern taiwan earthquakes and tectonic implications. *Terrestrial Atmospheric and Oceanic Sciences* 15, 647–666.
- Lee, J.C., Angelier, J., Chu, H.T., Hu, J.C., Jeng, F.S., 2001. Continuous monitoring of an active fault in a plate suture zone: a creepmeter study of the chihshang fault, eastern taiwan. *Tectonophysics* 333, 219–240.
- Lee, J.C., Angelier, J., Chu, H.T., Hu, J.C., Jeng, F.S., 2005. Monitoring active fault creep as a tool in seismic hazard mitigation. insights from creepmeter study at chihshang, taiwan. *Comptes Rendus Geoscience* 337, 1200–1207.
- Lee, J.C., Angelier, J., Chu, H.T., Yu, S.B., Hu, J.C., 1998. Plate-boundary strain partitioning along the sinistral collision suture of the philippine and eurasian plates: Analysis of geodetic data and geological observation in southeastern taiwan. *Tectonics* 17, 159–181.
- Lee, J.C., Fu-Shu, J., Hao-Tsu, C., Angelier, J., Jyr-Ching, H., 2000. A rod-type creepmeter for measurement of displacement in active fault zone. *Earth, Planets and Space* 52, 321–328.

- Lee, Y.H., Chen, G.T., Rau, R.J., Ching, K.E., 2008. Coseismic displacement and tectonic implication of 1951 longitudinal valley earthquake sequence, eastern taiwan. *Journal of Geophysical Research-Solid Earth* 113, 13.
- Mozziconacci, L., Delouis, B., Angelier, J., Hu, J.C., Huang, B.S., 2009. Slip distribution on a thrust fault at a plate boundary: the 2003 chengkung earthquake, taiwan. *Geophysical Journal International* 177, 609–623.
- Peyret, M., Dominguez, S., Cattin, R., Champenois, J., Leroy, M., Zajac, A., 2011. Present-day interseismic surface deformation along the longitudinal valley, eastern taiwan, from a ps-insar analysis of the ers satellite archives. *Journal of Geophysical Research* 116, B03402.
- Shyu, J.B.H., Chung, L.H., Chen, Y.G., Lee, J.C., Sieh, K., 2007. Re-evaluation of the surface ruptures of the november 1951 earthquake series in eastern taiwan, and its neotectonic implications. *Journal of Asian Earth Sciences* 31, 317–331.
- Thomas, M.Y., Avouac, J.P., Champenois, J., Lee, J.C., 2014. Spatio-temporal evolution of seismic and aseismic slip on the longitudinal valley fault, taiwan. *Journal of Geophysical Research*, in revision .
- Wu, Y.M., Chen, C.C., Zhao, L., Chang, C.H., 2008. Seismicity characteristics before the 2003 chengkung, taiwan, earthquake. *Tectonophysics* 457, 177–182.
- Wu, Y.M., Chen, Y.G., Shin, T.C., Kuochen, H., Hou, C.S., Hu, J.C., Chang, C.H., Wu, C.F., Teng, T.L., 2006. Coseismic versus interseismic ground deformations, fault rupture inversion and segmentation revealed by 2003 mw 6.8 chengkung earthquake in eastern taiwan. *Geophysical Research Letters* 33, L02312.